

Table 6

**Overview of Samples Used by Rind et al. (1998) to Determine Effect of Consent on CSA-Adjustment Relations in Male College Students**

Study	Contact <sup>a</sup>	Response <sup>b</sup>	Definition of CSA <sup>c</sup>	Dependent	Prevalence and effect sizes <sup>e</sup>		
				Measures <sup>d</sup>	N	CSA	r <sub>c</sub>
<b><u>Samples which specified CSA had to be unwanted</u></b>							
Bendixen et al, 1994	B	U	<18	Complaints (IA)	486	03%	.26
Collings, 1995	B	U	<18	BSI	284	29%	.29
Rau, 1994	B	U	<12 w/ 5+; 12-16 w/ 10+	PFQ, HSF-MMPI, FSSS	60	na	.10
Rew et al, 1991	B	U	<18 w/ Older partner	SES; JCS; GWBS	160	22%	.16
<b><u>"All types" (assumed by Rind et al. to contain wanted experiences)</u></b>							
Finkelhor, 1979, 1984	B	M	<13 w/ ≥16; 13-16 w/ 10+ (relative or unwanted)	FSSS	260	07%	.23
Fishman, 1991	B	M	<13 w/ 5+; 13-16 w/ 10+ or unwanted	SSD	148	21%	.12
Fromuthz & Burkhardt, 1989 [MW]	B	A	<13 w/ 5+ or ≥ 16; 13-16 w/ 10+	SCL-90; BDI-SF; LC; RSES	249	15%	.18
Fromuthz & Burkhardt, 1989 [SE]	B	A	<13 w/ 5+ or ≥ 16; 13-16 w/ 10+	SCL-90; BDI-SF; LC; RSES	324	13%	.01
Greenwald, 1994	B	M	<16 w/ 5+; ≥ 16 w/ 10+; unwanted; intrafamilial	RSSS	111	09%	-.22
Hatfield, 1988	C	A	<14 w/ 5+	HSCL-Mod	213	12%	.14
Predieri, 1992	B	A	<13 w/ 5+; 13-16 w/8+; force; authority figure	DSFI, MMPI Sc5	62	na	.09
Preuss, 1988	B	A	<13 w/ 5+; 13-17 w/10+; force	MAST; bulimia (IA)	288	20%	.09
Sarbo, 1985	B	M	<12 w/ ≥16; <16 w/ relative ≥16; unwanted	CAQ	112	24%	.12
Urquiza, 1989	C	A	<18 w/ 5+	SC, TSSS, MAST	88	na	.19

**Note.** <sup>a</sup> C = contact only; B = both contact and noncontact. <sup>b</sup> U = only unwanted sex included in the definition; A = all types of sexual experiences, unwanted and wanted; M = mixed, unwanted experiences are included in the definition for older age groups. <sup>c</sup> Upper age of "child" is given first; then, minimum age of perpetrator is given (e.g., w/ 5+ means with someone at least 5 years older); last, other condition for CSA are given. <sup>d</sup> Objective measures included in effect size estimates: BDI = Beck Depression Inventory-Short Form (Beck, Ward, Mendelson, Mock & Erbaugh, 1961); BSI = Brief Symptom Inventory (Derogatis & Spencer, 1982); CAQ = Clinical Analysis Questionnaire (Cattell, 1971); DSFI = Derogatis Sexual Functioning Inventory (Derogatis & Melisaratos, 1979); FSSS = Finkelhor Sexual Self-Esteem Scale (Finkelhor, 1984); GWBS = General Well-Being Schedule (McDowell & Newell, 1987); HSCL = Hopkins Symptom Check List (Derogatis et al., 1974); HSCL-Mod = Hopkins Symptom Check List - Modified (Briere & Runtz, 1985; Derogatis et al., 1974); HSF-MMPI = Hugo Short Form of the MMPI (Hugo, 1971); IA = investigator-authored item; LOCS = Locus of Control Scale (Coleman et al., 1966); MAST = Michigan Alcoholism Screening Test (Brady, Foulks, Childress, & Pertschuk, 1982); MMPI = Minnesota Multiphasic Personality Inventory (Hathaway & McKinley, 1967); PSQ = Psychosexual Functioning Questionnaire (Schover, Friedman, Weiler, Heinman, & LoPiccolo, 1982); RSES = Rosenberg Self-Esteem Scale (Rosenberg, 1965); RSSS = Romantic and Sexual Self-esteem Scale (Reed, 1987); SC = Symptom Checklist (Briere, 1984); SCL-90 = Symptom Checklist (Derogatis, Lipman, & Covi, 1973); SES = Self-Efficacy Scale (Sherer et al., 1982); SSD = Sexual Self-Esteem and Dysfunction (investigator authored); TSSS = Tennessee Self-Concept Scale (Fitts, 1965); Note: subjective perceptions were not included. <sup>e</sup> N<sub>s</sub> are number of subjects in analyses of psychological correlates of CSA; effect sizes are at the sample-level; r<sub>s</sub> are point-biserial effect sizes corrected for base rate differences of CSA, but not dichotomization.